



Accession # 00385382
Female Patient



Adrenal


Ordering Physician:
 Regenerus Laboratories Ltd

DOB: 1981-06-30
Age: 38
Gender: Female

Last Menstrual Period:

2019-08-04
Collection Times:
 2019-08-27 08:15AM
 2019-08-27 10:20AM
 2019-08-26 05:15PM
 2019-08-26 10:00PM

Category	Test		Result	Units	Normal Range
Creatinine (Urine)					
	Creatinine A (Waking)	Within range	0.82	mg/ml	0.2 - 2
	Creatinine B (Morning)	Within range	1.13	mg/ml	0.2 - 2
	Creatinine C (Afternoon)	Within range	0.76	mg/ml	0.2 - 2
	Creatinine D (Night)	Within range	1.0	mg/ml	0.2 - 2
Daily Free Cortisol and Cortisone (Urine)					
	Cortisol A (Waking)	Above range	53.7	ng/mg	10 - 50
	Cortisol B (Morning)	Above range	204.0	ng/mg	30 - 130
	Cortisol C (Afternoon)	High end of range	26.8	ng/mg	7 - 30
	Cortisol D (Night)	Within range	8.9	ng/mg	0 - 14
	Cortisone A (Waking)	High end of range	105.4	ng/mg	40 - 120
	Cortisone B (Morning)	Above range	232.7	ng/mg	90 - 230
	Cortisone C (Afternoon)	Within range	89.0	ng/mg	32 - 110
	Cortisone D (Night)	Within range	26.1	ng/mg	0 - 55
	24hr Free Cortisol	Above range	293.3	ng/mg	65 - 200
	24hr Free Cortisone	Above range	453.2	ng/mg	220 - 450
Cortisol Metabolites and DHEA-S (Urine)					
	a-Tetrahydrocortisol (a-THF)	Above range	502.0	ng/mg	75 - 370
	b-Tetrahydrocortisol (b-THF)	Within range	1792.0	ng/mg	1050 - 2500
	b-Tetrahydrocortisone (b-THE)	Within range	3317.0	ng/mg	1550 - 3800
	Metabolized Cortisol (THF+THE)	Within range	5611.0	ng/mg	2750 - 6500
	DHEA-S	Within range	241.0	ng/mg	20 - 750




Stress (or inflammation) causes the brain to release ACTH, which stimulates the adrenal glands to make hormones

DHEA-S Ranges

Age	Range
20-39	60-750
40-60	30-350
>60	20-150







Metabolized Cortisol (THF+THE)
(Total Cortisol Production)




More cortisone metabolites (THE) More cortisol metabolites (THE)

NOTE: This 11b-HSD index measures the balance of cortisol and cortisone metabolites which best reflects the overall balance of active cortisol and inactive cortisone systemically.



24hr Free Cortisone
(A+B+C+D)

Cortisol and Cortisone interconvert (11b-HSD)






24hr Free Cortisol
(A+B+C+D)

The first value reported (Waking "A") for cortisol is intended to represent the "overnight" period. When patients sleep through the night, they collect just one sample. In this case, the patient did not report waking up during the night to collect a sample, so the "Waking (A)" cortisol and cortisone values should accurately represent the entirety of the overnight period.


Provider Notes

How to read the DUTCH report

This report is not intended to treat, cure or diagnose any specific diseases. The graphic dutch dials in this report are intended for quick and easy evaluation of which hormones are out of range. Results below the left star are shaded yellow and are below range (left). Results between the stars and shaded green are within the reference range (middle). Results beyond the second star and shaded red are above the reference range (right). Some of these hormones also change with age, and the age-dependent ranges provided should also be considered.



In a few places on the graphical pages, you will see fan-style gauges. For sex hormones, you will see one for the balance between 5a/5b metabolism as well as methylation. For adrenal hormones, you will see one to represent the balance between cortisol and cortisone metabolites. These indexes simply look at the ratio of hormones for a preference. An average or "normal" ratio between the two metabolites (or groups of metabolites) will give a result in the middle (as shown here). If the ratio between the metabolites measured is "low" the gauge will lean to the left and similarly to the right if the ratio is higher than normal.



Patient or Sample Comments

Throughout the provider comments you may find some comments specific to your situation or results. These comments will be found in this section or within another section as appropriate. Comments in other sections that are specific to your case will be in **bold**.

The following video link(s) may help those new to dutch testing to understand the results. If you only have a hardcopy of the results, the video names can be easily found in our video library at www.DutchTest.com. Be aware that our reporting format has recently undergone some cosmetic changes, so the results on the video may look slightly different. These results and videos are NOT intended to diagnose or treat specific disease states.

This video may assist with the interpretation of the Adrenal (cortisol) results: [Cortisol tutorial video](#)

The patient reports regular menstrual cycles.

The patient reports symptoms of both estrogen deficiency and excess. Levels of estrogen (and metabolites) should be carefully reviewed.

The patient reports significant symptoms of both androgen deficiency and excess. Results and symptoms should be reviewed carefully.

The patient reported significant fatigue in the afternoon/evening, but not in the morning.

DUTCH Adrenal

The HPA-Axis refers to the communication and interaction between the hypothalamus (H) and pituitary (P) in the brain down to the adrenal glands (A) that sit on top of your kidneys. When a physical or psychological stressor occurs, the hypothalamus tells the pituitary to make ACTH, a hormone. ACTH stimulates the adrenal glands to make the stress hormone, cortisol and to a lesser extent DHEA and DHEA-S. Normally, the HPA-axis production follows a daily pattern in which cortisol rises rather rapidly in the first 10-30 minutes after waking in order to help with energy, then gradually decreases throughout the day so that it is low at night for sleep. The cycle starts over the next morning. Abnormally high activity occurs in Cushing's Disease where the HPA-axis is hyper-stimulated causing cortisol to be elevated all day. The opposite is known as Addison's Disease, where cortisol is abnormally low because it is not made appropriately in response to ACTH's stimulation. These two conditions are somewhat rare. Examples of more common conditions related to less severely abnormal cortisol levels include fatigue, depression, insomnia, fibromyalgia, anxiety, inflammation and more.

Only a fraction of cortisol is "free" and bioactive. This fraction of cortisol is very important, but levels of metabolized cortisol best represent overall production of cortisol therefore both should be taken into account to correctly assess adrenal function.

When evaluating cortisol levels, it is important to assess the following:

- **The overall up-and-down pattern of free cortisol throughout the day, looking for low and high levels:** Abnormal results should be considered along with related symptoms. Remember that with urine results, the "waking" sample reflects the night's total for free cortisol. The sample collected two hours after waking captures the cortisol awakening response, which is typically the time with the most cortisol secretion.
- **The sum of the free cortisol as an expression of the overall tissue cortisol exposure:**

This total of four free cortisol measurements is the best way to assess the total of free cortisol throughout the day, and this result correlates reasonably well to a true 24-hour urine free cortisol. Do be aware that this measurement does not take into account transitory shifts in cortisol in the late morning or early afternoon.

• **The total level of cortisol metabolites:**

We call this calculation "Metabolized Cortisol" which is the sum of a-THF, b-THF and b-THE (the most abundant cortisol metabolites). While free cortisol is the best assessment for tissue levels of cortisol, it only represents 1-3% of the total produced. The majority of cortisol results in a urine metabolite and the total of these metabolites best represents the total glandular output of cortisol for the day. When overall production is much higher than free cortisol levels, cortisol clearance may be increased (as seen in hyperthyroidism, obesity, etc.) The most common reason for sluggish cortisol clearance (assumed when free cortisol levels are much higher than metabolized cortisol) is low thyroid.

Please note, cortisol and cortisone ranges have been updated as of 8.20.2019. These slight changes represent a range optimization following a methodological change.

Overall free cortisol levels are elevated, but metabolized cortisol (the best marker for overall cortisol production) is within range. Cortisol clearance may be a bit sluggish, which keeps free cortisol levels elevated in spite of normal overall production.

• **A potential preference for cortisol or cortisone (the inactive form):**

Looking at the comparison between the total for free cortisol and free cortisone is NOT the best indication of a person's preference for cortisol or cortisone. The kidney converts cortisol to cortisone in the local tissue. This localized conversion can be seen by comparing cortisol (free) and cortisone levels. To see the patient's preference systemically, it is best to look at which *metabolite* predominates (THF or THE). This preference can be seen in the fan style gauge. This is known as the 11b-HSD index. The enzyme 11b-HSD II converts cortisol to cortisone in the kidneys, saliva gland and colon. 11b-HSD I is more active in the liver, fat cells and the periphery and is responsible for reactivating cortisone to cortisol. Both are then metabolized by 5a-reductase to become tetrahydrocortisol (THF) and tetrahydrocortisone (THE) respectively.

The patient's THF/THE ratio implies a preference for cortisone (relative to cortisol). Because cortisol levels are not low, this may have some beneficial effect in keeping cortisol levels from being even higher than reported here. The patient does report some fatigue so the overall case must be considered carefully in terms of how to best address symptoms of fatigue as it may not be due to dysfunctional HPA-Axis if the diurnal pattern is relatively normal. If the diurnal pattern is normal, consider other causes of fatigue such as anemia among others.

Urine Hormone Testing - General Information

What is actually measured in urine? In blood, most hormones are bound to binding proteins. A small fraction of the total hormone levels are "free" and unbound such that they are active hormones. These free hormones are not found readily in urine except for cortisol and cortisone (because they are much more water soluble than, for example, testosterone). As such, free cortisol and cortisone can be measured in urine and it is this measurement that nearly all urinary cortisol research is based upon. In the DUTCH Adrenal Profile the diurnal patterns of free cortisol and cortisone are measured by LC-MS/MS.

All other hormones measured (cortisol metabolites, DHEA, and all sex hormones) are excreted in urine predominately after the addition of a glucuronide or sulfate group (to increase water solubility for excretion). As an example, Tajic (Natural Sciences, 1968 publication) found that of the testosterone found in urine, 57-80% was testosterone-glucuronide, 14-42% was testosterone-sulfate, and negligible amounts (<1% for most) was free testosterone. The most likely source of free sex hormones in urine is from contamination from hormonal supplements. To eliminate this potential, we remove free hormones from conjugates (our testing can be used even if vaginal hormones have been given). The glucuronides and sulfates are then broken off of the parent hormones, and the measurement is made. These measurements reflect the bioavailable amount of hormone in most cases as it is only the free, nonprotein-bound fraction in blood/tissue that is available for phase II metabolism (glucuronidation and sulfation) and subsequent urine excretion.

Disclaimer: the filter paper used for sample collection is designed for blood collection, so it is technically considered "research only" for urine collection. Its proper use for urine collection has been thoroughly validated.

